

Clarus[®]
Isotretinoin Capsules, USP

MONTHLY PATIENT REMINDER SLIPS (for female patients)



DON'T FORGET

- You **MUST** have a **NEGATIVE** pregnancy test to continue treatment with CLARUS[®]
- You **MUST** abstain from sexual intercourse

OR

Use **TWO** reliable methods of Birth Control at the same time during your CLARUS[®] treatment;

AND

For at least **ONE** month after completion of your CLARUS[®] treatment

- IF** you become or **THINK** you could be pregnant, consult your Doctor immediately and stop taking CLARUS[®].

QUESTIONS?

To receive confidential contraception counselling from a Health Care Professional, please call: **1-877-776-7711**

OFFICE STAMP

DR.'S
INITIALS

DON'T FORGET

- You **MUST** have a **NEGATIVE** pregnancy test to continue treatment with CLARUS[®]
- You **MUST** abstain from sexual intercourse

OR

Use **TWO** reliable methods of Birth Control at the same time during your CLARUS[®] treatment;

AND

For at least **ONE** month after completion of your CLARUS[®] treatment

- IF** you become or **THINK** you could be pregnant, consult your Doctor immediately and stop taking CLARUS[®].

QUESTIONS?

To receive confidential contraception counselling from a Health Care Professional, please call: **1-877-776-7711**

OFFICE STAMP

DR.'S
INITIALS

DON'T FORGET

- You **MUST** have a **NEGATIVE** pregnancy test to continue treatment with CLARUS[®]
- You **MUST** abstain from sexual intercourse

OR

Use **TWO** reliable methods of Birth Control at the same time during your CLARUS[®] treatment;

AND

For at least **ONE** month after completion of your CLARUS[®] treatment

- IF** you become or **THINK** you could be pregnant, consult your Doctor immediately and stop taking CLARUS[®].

QUESTIONS?

To receive confidential contraception counselling from a Health Care Professional, please call: **1-877-776-7711**

OFFICE STAMP

DR.'S
INITIALS

DON'T FORGET

- You **MUST** have a **NEGATIVE** pregnancy test to continue treatment with CLARUS[®]
- You **MUST** abstain from sexual intercourse

OR

Use **TWO** reliable methods of Birth Control at the same time during your CLARUS[®] treatment;

AND

For at least **ONE** month after completion of your CLARUS[®] treatment

- IF** you become or **THINK** you could be pregnant, consult your Doctor immediately and stop taking CLARUS[®].

QUESTIONS?

To receive confidential contraception counselling from a Health Care Professional, please call: **1-877-776-7711**

OFFICE STAMP

DR.'S
INITIALS